

(43) International Publication Date 30 November 2000 (30.11.2000)

PCT

(10) International Publication Number WO 00/71116 A1

C07D 207/34

M., Dileep [IN/IN]; Vaishali Apartments, Pocket A-12, 3C, Kalkaji Extension, New Delhi 110 019, Maharashtra (IN).

(21) International Application Number: PCT/IB00/00014

(22) International Filing Date: 6 January 2000 (06.01,2000)

A61K 31/40,

•

(51) International Patent Classification7:

English

(26) Publication Language:

English

(30) Priority Data:

775/DEL/99

(25) Filing Language:

25 May 1999 (25.05.1999) IN

(71) Applicant (for all designated States except US): RAN-BAXY LABORATORIES LIMITED [IN/IN]; 19 Nehru Place, New Delhi 110 019, Maharashtra (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KUMAR, Yatendra [IN/IN]; U-26/5, DLF Qutab Enclave, Phase - III, Gurgaon 122 001, Haryana (IN). THAPER, Rajesh, Kumar [IN/IN]; B-26, Sunshine Apartments, Block "C", Sushant Lok - I, Gurgaon 122 002, Haryana (IN). KUMAR, S.,

(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

1/11116 A

(57) Abstract: A process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises: (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent; (b) adding a non-polar hydrocarbon anti-solvent or adding the dissolved atorvastatin to the non-polar anti-solvent to precipitate out atorvastatin calcium; and (c) removing the solvent by filtration to afford amorphous atorvastatin calcium.

PROCESS FOR THE PRODUCTION OF AMORPHOUS ATORVASTATIN CALCIUM

5

FIELD OF THE INVENTION

The present invention relates to a process for the production of amorphous atorvastatin calcium.

10

15

20

BACKGROUND OF THE INVENTION

Atorvastatin is chemically [R-(R*,R*)]-2-(4-fluoro-phenyl)-β dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H pyrrole-1-heptanoic acid. Atorvastatin calcium, a synthetic HMG-CoA reductase inhibitor, is used for the treatment of hyperlipidemia and hypercholesterolemia, both of which are risk factors for arteriosclerosis and coronary heart disease. Open dihydroxy carboxylic acid, lactone and various salt forms of atorvastatin have been synthesized.

United States Patent 5,273,995, describes that R-form of the ring opened acid form has surprising inhibition of the biosynthesis of cholesterol. Atorvastatin in its calcium salt form, i.e. $[R-(R^*,R^*)]-2-(4-fluoro-phenyl)-\beta$, δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbomyl]-1H-pyrrole-1-heptanoic acid calcium salt (2:1) having Formula 1:

5

10

15

20

is more suited to formulations and has been recommended as a drug.

United States patents 5,003,080; 5,097,045; 5,103,024; 5,124,482; 5,149,837; 5,248,793; 5,280,126; 5,342,952, which are herein incorporated by reference, describe various processes and key intermediates for preparing atorvastatin.

Atorvastatin calcium produced by the processes described in the above mentioned United States patents does not give amorphous atorvastatin consistently but gives a mixture of its crystalline and amorphous forms, which has unsuitable filtration and drying characteristics and are not suitable for large-scale production.

PCT application, WO 97/03959, discloses novel crystalline forms of atorvastatin calcium designated as Form I, Form II, and Form IV and method for their preparation which provide more favourable filtration and drying characteristics.

PCT application WO 97/03960 describes a procedure for converting the crystalline form of atorvastatin to the amorphous form. Process disclosed therein comprises dissolving crystalline form-I atorvastatin in a non-hydroxylic solvent like tetrahydrofuran or mixtures of tetrahydrofuran and toluene. The process involves complete removal of the solvent under high temperature (about 90°C) and high vacuum (about 5mm) using capital intensive equipment. Exposure of the material to high temperature for several days leads to degradation of the product. This makes the process very inconvenient to operate at a large scale. Slow removal of solvents at a manufacturing scale renders this process as inefficient cost-wise and less productive.

10

5

SUMMARY OF THE INVENTION

It is an objective of the present invention to provide an efficient process for the production of amorphous atorvastatin, which eliminates the problems of prior art and is convenient to operate on a commercial scale.

15

Accordingly, the present invention provides a process for the preparation of atorvastatin calcium in an amorphous form which comprises dissolving crystalline atorvastatin in a non-hydroxylic solvent, adding a suitable non-polar hydro-carbon solvent and recovering atorvastatin from a solution thereof, by solvent precipitation, isolating and drying the product.

20

Generally, the product can be isolated by any standard method known in the art such as by filtration, centrifugation or decantation. Typically, this product is isolated by filtration when any of the solvents within the scope of the process are used.

Major advantages of the present invention compared to the prior art processes are:

i. elimination of the need to remove solvent by drying techniques.

- ii. less time consuming with improved filtration.
- iii. easy to operate on large-scale.
- iv. reproducibly produces amorphous product having allowable levels of residual solvents.

The present invention thus provides a novel process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:

- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent to precipitate out the material; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium

The non-hydroxylic solvent is selected from a group of solvents, which have the ability to dissolve crystalline atorvastatin and includes tetrahydrofuran. Suitable non-polar hydrocarbon solvents are selected from a group consisting of: n-hexane, n-heptane, cyclohexane, hexane fraction, heptane fraction or the like. In a preferred embodiment of this invention, the non-hydroxylic solvent is tetrahydrofuran and antisolvent is n-hexane, cyclohexane or n-heptane.

Generally, crystalline atorvastatin calcium is dissolved in a non-hydroxylic solvent, e.g. tetrahydrofuran, at a concentration of about 15% w/v to about 40% w/v, preferably at a concentration of about 25% w/v to about 15% w/v at ambient

10

5

15

20

temperature and a non-polar hydrocarbon, preferably n-hexane, cyclohexane or n-heptane, is added at 0°C to 50°C, preferably at 20°C to 25°C. The product is recovered by filtration at ambient temperature. Filtration, which is fast and smooth, is carried out using nutsche filtration or centrifuge filtration. Preferably, nutsche filtration is used on large scale preparation. Filtered material, a semi-dry powder, is further dried to remove surface solvents in a vacuum tray drier, tray drier, fluid bed drier or a rotary vacuum drier to afford amorphous material. Preferably, material is dried in a vacuum tray drier at about 20°C to about 80°C for 6 hours to 24 hours. Most preferably, drying is carried out at about 50°C to about 60°C for 12 hours.

10

5

Quantity of antisolvent varies from 5 times to 50 times the input of crystalline atorvastatin calcium depending upon its solution in non-hydroxylic solvent. Preferably, the quantity of antisolvent used is about 20 times to about 40 times the input of crystalline atorvastatin calcium to make overall concentration of about 5% w/v to about 2.5 w/v%.

15

20

Amorphous atorvastatin calcium prepared according to the process of the present invention may be characterized by its x-ray powder diffration pattern (Figures 2) as shown in the accompanied drawings. X-ray powder diffration patterns (Figures 2) show no peaks which are characteristic of a crystalline atorvastatin calcium (Figure 1 of the accompanied drawings) thus demonstrating the amorphous nature of the product.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is the diffractogram of crystalline atorvastatin calcium. The horizontal

axis represents 2θ and the vertical axis corresponds to peak intensity.

Figure 2 is diffractogram of amorphous atorvastatin calcium. The horizontal

axis represents 2θ and the vertical axis corresponds to peak intensity.

The present invention is illustrated by the following examples, which are not intended to limit the effective scope of the claims.

DETAILED DESCRIPTION OF THE INVENTION

Example 1

 $[R-(R^*,R^*)-2-(4-fluorophenyl)-\beta, \qquad \delta-dihydroxy-5-(1-methylethyl)-3-phenyl-4-\\ [(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium).$

Method A

5

10

15

20

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. Clear solution so obtained was added slowly to cyclohexane (350 lt) under nitrogen atmosphere. It was vigorously stirred maintaining temperature at 20-25°C. The precipitated product was centrifuged and dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.5 kg) in an

amorphous form was obtained having residual solvent levels of 0.01% w/w tetrahydrofuran and 0.6% w/w cyclohexane. X-ray powder diffraction pattern (Figure 2 as shown in the accompanied drawings) demonstrate the amorphous nature of the product.

5

Method B

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. To a clear solution of atorvastatin, cyclohexane (350 lt) was added under vigorous stirring at 20 to 25°C. The precipitated mass was further stirred for 30 minutes and filtered in a centrifuge. The product was dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.6 kg) in an amorphous form was obtained having residual solvent levels of 0.01% w/w for tetrahydrofuran and 0.7% w/w for cyclohexane. X-ray powder diffraction pattern demonstrates the amorphous nature of the product.

15

10

Example 2

20

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-hexane instead of cyclohexane

to give amorphous atorvastatin (9.5 kg.). X-ray crystallography confirmed the amorphous nature of the product.

Example 3

 $[R-(R^*,R^*)]-2-4-fluorophenyl)-\beta, \qquad \delta-dihydroxy-5-(1-methylethyl)-3-phenyl-4-\\ [(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium)$

10

15

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-heptane instead of cylcohexane to give amorphous atorvastatin (9.6 kg). X-ray crystallography examination confirmed the amorphous nature of the product.


While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.


WE CLAIM:

 A process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:

- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent or adding the dissolved atorvastatin to the non-polar anti-solvent to precipitate out atorvastatin calcium; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium.
- 2. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is chosen from a group of non-polar hydrocarbon solvents comprising n-hexane, cyclohexane or n-heptane.
- 3. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-hexane.
- 4. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is cylcohexane.
- 5. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-heptane.
- 6. The process of claim 1, wherein said amorphous atorvastatin calcium is isolated by filtration.







Interr nal Application No PCT/IB 00/00014

A 61 455-	TO A TION OF CHIP ISCT MATTER		
IPC 7	A61K31/40 C07D207/34		
According to	International Patent Classification (IPC) or to both national class	ification and IPC	
B. FIELDS	SEARCHED		
IPC 7			
	tion searched other than minimum documentation to the extent th		
Electronic d	eta base consulted during the international search (name of data	a base and, where practical, search terms used)	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the	e relevant passages	Relevant to claim No.
A	WO 97 03959 A (WARNER LAMBERT (CHRISTOPHER A (US); JENNINGS RI 6 February 1997 (1997-02-06) cited in the application		1-6
A	DE 33 27 449 A (GLAXO GROUP LTG 2 February 1984 (1984-02-02) page 9, line 7 - line 16 page 12, line 10 -page 13, line	1-6	
A	WO 97 03960 A (WARNER LAMBERT (US); SCHWEISS DIETER (US)) 6 February 1997 (1997-02-06) cited in the application claim 1	CO ;LIN MIN	1-6
		-/	
X Fu	rther documents are listed in the continuation of box C.	Patent family members are listed	I in annex.
* Special	categories of cited documents:	"T" later document published after the int	emational filing date
cons	ment defining the general state of the art which is not sidered to be of particular relevance	or priority date and not in conflict with cited to understand the principle or the invention	n the application but
	r document but published on or after the international grate	"X" document of particular relevance; the cannot be considered novel or canno	nt be considered to
"L" docur which	nent which may throw doubts on priority claim(s) or this cited to establish the publication date of another tion or other special reason (as specified)	involve an inventive step when the d "Y" document of particular relevance; the cannot be considered to involve an in	ocument is taken alone claimed invention
othe	ment referring to an oral disclosure, use, exhibition or or means	document is combined with one or n ments, such combination being obvi in the art.	ore other such docu-
	ment published prior to the international filing date but r than the priority date claimed	"&" document member of the same pater	
Date of the	ne actual completion of the international search	Date of mailing of the international se	earch report
	10 May 2000	17/05/2000	
Name an	d mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2	Authorized officer	
	NL – 2280 HV Rijswijk Tel. (+31-70) 340–2040, Tx. 31 651 epo nl, Fax: (+31-70) 340–3016	De Jong, B	

1

Intern 1al Application No PCT/IB 00/00014

	tion) DOCUMENTS CONSIDERED TO BE RELEVANT	In.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
1	WO 97 03958 A (WARNER LAMBERT CO ;MCKENZIE ANN T (US)) 6 February 1997 (1997-02-06) cited in the application	1-6

1

information on patent family members

inten nal Application No PCT/IB 00/00014

					1/10	10/00014
Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 9703959	A	06-02-1997	AU BG	6484296 102187	A	18-02-1997 30-10-1998
			BR	9609872		23-03-1999
			CA CN	2220018 1190955		06-02-1997 19-08-1998
			CZ	9800121		14-10-1998
			EP	0848705		24-06-1998
			HR	960339		30-04-1998
			HU	9900678		28-07-1999
			IL	122118		14-07-1999
			JP	11509230		17-08-1999
			NO	980207		16-01-1998 25-05-1998
			PL SK	324496 6298		07-10-1998
			US	5969156		19-10-1999
DE 3327449	A	02-02-1984	AT	382154 276783		26-01-1987 15-06-1986
			AT AU	566881		05-11-1987
			AU	1741783		02-02-1984
			BE	897422		30-01-1984
			CA	1240313		09-08-1988
			CH	657134		15-08-1986
			CZ	280528		14-02-1996 15-03-1988
			CS CY	8305687 1434		02-09-1988
			DE	3374010		12-11-1987
			ĎK	68392		25-05-1992
			DK	349083		31-01-1984
,			EP	0107276		02-05-1984
			ES ES	524590 8505689		01-06-1985 01-10-1985
			FI	832757		31-01-1984
			FŘ	2531087		03-02-1984
			GB	2127401		11-04-1984
			GR	79349		22-10-1984
			HK	84288		28-10-1988 29-09-1986
			HU IE	190603 55748		02-01-1991
			IL	69375		31-12-1986
1			IT	1168206		20-05-1987
			JP	2025666		26-02-1996
			JP	7030084		05-04-1995
			JP KE	59044391 3805		12-03-1984 03-06-1988
			KR	9100046		19-01-1991
			LÜ	84935		23-11-1983
			MY	5887	' A	31-12-1987
			NL	8302705		16-02-1984
			NO NZ	832773		31-01-1984 14-03-1986
			NZ Pl	205083 243228		27 - 08-1984
			PT	77135		01-08-1983
			SE	453195		18-01-1988
			SE	8304208	3 A	31-01-1984
			SG	26088		15-07-1988
			SI SK	8311558 403191		31 - 12-1995 11 - 07-1995
			2K	40319	L M	11 -01 1333

information on patent family members

Inten nal Application No PCT/IB 00/00014

Patent document cited in search report		Publication date		atent family nember(s)	Publication date
DE 3327449	Α	<u> </u>	SU	1266471 A	23-10-1986
			US	4994567 A	19-02-1991
			US	5013833 A	07-05-1991
			US	4562181 A	31-12-1985
			US	4820833 A	11-04-1989
			YU	155883 A	30-04-1986
			ZA	8305579 A	26-09-1984
			ZW	17383 A	26-10-1983
WO 9703960	A	06-02-1997	AU	700794 B	14-01-1999
			AU	6497896 A	18-02-1997
			BG	102188 A	31-08-1998
			BR	9609714 A	23-02-1999
			CA	2220455 A	06-02-1997
			CN	1190956 A	19-08-1998
			CZ	9800122 A	16-12-1998
			EP	0839132 A	06-05-1998
			HR	960312 A	28-02-1998
			IL	122161 A	14-07-1999
			JP	11510486 T	14-09-1999
			NO	980209 A	16-01-1998
			PL	324463 A	25-05-1998
			SK	5898 A	05-08-1998
WO 9703958	Α	06-02-1997	AU	6484196 A	18-02-1997
			BG	102186 A	30-10-1998
			BR	9610567 A	06-07-1999
			CA	2220458 A	06-02-1997
			CN	1190957 A	19-08-1998
			CZ	9800123 A	17-06-1998
			EP	0848704 A	24-06-1998
			HR	960313 A	30-04-1998
			HN	9901687 A	28-10-1999
			IL	122162 A	14-07-1999
			JP	11509229 T	17-08-1999
			NO Pl	980208 A 324532 A	16-01-1998 08-06-1998
			SK	324532 A 5998 A	06-05-1998
			3K	DYYO A	00-05-1996